GOAT AFFECTED WITH MANGE (Sarcoptes spp.)

How To Diagnose:



Severe exfoliative skin lesions over the muzzle, ears, around the eyes and other parts of the body.



In kids, there will be depression and stunted growths if affected during early age.



Without appropriate treatment, lesions will spread throughout the body and in severe infection, will result in mortality.



Intense pruritis, scaling of different body parts. Loss of hair from the affected body parts.

Whom To Contact:

- Local Veterinary Officer/ Field Assistant.
- ADMaC, Core Lab- I, AAU, Khanapara
 - Dr. S. K. Das, PI, ADMaC.
 E mail drskdas53@gmail.com, Mob No. 09435014705
 - ◆ Dr. N.N.Barman, Co PI, ADMaC.
 E mail nnbarman@gmail.com, Mob No.0943555 8788

What will you do:

- Do not allow goats to move in market places.
- Separate the affected ones from the healthy and keep them in clean hygienic sheds/places.
- Persons should take protective measures while attending the affected goat.
- Inform nearest Veterinary Doctor/Field Assistant.
- Send appropriate samples through Veterinary Doctor/Field Assistant to the nearest diagnostic laboratory/ ADMaC Core Lab-I, Khanapara for confirmation of the disease.
- Burry dead goats in 5 to 6 feet deep pit putting sufficient amount of common salt/lime or washing soda.
- Affected animals can be treated at fever phase.